

The chemical potential of the electron gas on a one dimensional  
lattice

**Vladan Čelebonović**

Institute of Physics, P.O.B. 57, 11001 Beograd, Yugoslavia

Abstract: The chemical potential of the electron gas on a one-dimensional lattice is determined within the discrete Hubbard model. The result will have applications in theoretical studies of transport properties of quasi one-dimensional organic conductors such as the Bechgaard salts.

## Introduction

Quasi one-dimensional (Q1D) organic metals were discovered in 1980 [1], [2], [2a]. It was soon shown [3] that the electrical conductivity of these materials could not be described by the theory of electrical conductivity of normal metals. The Q1D organic metals are, (since [4]), considered within the framework of the Hubbard model on a one-dimensional lattice. In spite of the fact that this model is one of the simplest in statistical mechanics, it is a subject of intensive theoretical studies. The model is solvable in one dimension, and the chemical potential is known only for the special case of a one-dimensional half-filled band, for which  $\mu = 0$  [5]. It is usually assumed that the electrons in Q1D organic metals form a normal Fermi liquid (however, see [6]), which means that

$$n_k = 1 / [1 + \exp \beta(\varepsilon_k - \mu)] \quad (1)$$

The electronic energies are given by  $\varepsilon_k = -2t\cos(ks)$  [5]. All the symbols have their usual meaning,  $t$  denotes the hopping and  $s$  the lattice constant.

The purpose of the present letter is to determine the dependence on the band-filling and temperature of the chemical potential of the electron gas on a one-dimensional lattice. A similar problem has been investigated in quantum field theory, where a special prescription had to be proposed for introducing the chemical potential in a lattice regulated field theory (see, for example, [15]). The knowledge of the function  $\mu = \mu(n, \beta, t)$  is of paramount importance for the calculation of the electrical conductivity and optical response functions of correlated electron systems [13], [14]. Deviating the filling from  $1/2$  is a theoretical procedure for representing the effects of doping on the material. Preliminary results of this calculation were recently published [5a].

The calculation which will be reported in this letter is a first step towards a theoretical study of the effects of doping on the transport properties and optical response functions of Q1D organic metals. For a recent example of an experimental study of these effects see [7]. Apart from the Q1D organic metals, the calculation to be reported is of interest for general one-dimensional models of solids (such as the Kronig-Penney model) [8].

## The calculation

The starting point for the determination of the chemical potential is the following relation (for example, [9]):

$$n = 2 \int 1 / [1 + \exp \beta(\varepsilon_k - \mu)] dp / (2\pi\hbar) \quad (2)$$

where all the symbols have their usual meaning,  $p = \hbar k$ , and  $n$  denotes the number of electrons per site ( $n = 1$  corresponds to filling  $1/2$ ). The limits of integration are  $\pm\pi\hbar/s$ . Using the fact that in the 1D Hubbard model  $\varepsilon_k = -2t\cos(ks)$ , it follows

$$dp = \hbar dk = (\hbar/s) (4t^2 - \varepsilon^2)^{-1/2} d\varepsilon \quad (3)$$

Inserting (3) into (2) one gets, after some algebra, that

$$n = 1/(\pi s) \int_{-2t}^{2t} 1 / [1 + \exp \beta(\varepsilon_k - \mu)] (4t^2 - \varepsilon^2)^{-1/2} d\varepsilon \quad (4)$$

and the problem is to find a solution of this integral equation. Instead of using the full rigour of the theory of integral equations (such as [10]), the problem can be solved by a suitable development of the Fermi function.

It has recently been shown [11] that the Fermi function can be represented in the following form:

$$1 / [1 + \exp \beta(\varepsilon_k - \mu)] = \Theta(\mu - \varepsilon) - \sum_{k=0}^{\infty} A_{2k+1} \beta^{-2(k+1)} \delta^{(2k+1)}(\varepsilon - \mu) \quad (5)$$

where  $A_{2k+1} = (-1)^{k+2} [2(2^{2k+1} - 1) \pi^{2k+2} / (2k+2)!] B_{2k+2}$

In eq. (5)  $\Theta$  denotes the step function,  $\delta^{(2k+1)}(\varepsilon - \mu)$  are the derivatives of the  $\delta$  function and  $B_{2k+2}$  are Bernoulli's numbers [12]. Inserting (5) into (4) it follows that

$$n = 1 / (\pi s) \int_{-2t}^{2t} [\Theta(\mu - \varepsilon) - \sum_{k=0}^{\infty} A_{2k+1} \beta^{-2(k+1)} \delta^{(2k+1)}(\varepsilon - \mu)] (4t^2 - \varepsilon^2)^{-1/2} d\varepsilon \quad (6)$$

This equation can be integrated by using the following relation, valid for an arbitrary function  $f(x)$  and its  $n$ -th order derivative:

$$\int \delta^{(n)}(x - x_0) f(x) dx = (-1)^n f^{(n)}(x_0) \quad (7)$$

Combining (7) and (6), and limiting (5) to terms with  $k \leq 2$  one finally obtains

$$1 / (\pi s) [\pi + (\pi^2/6\beta^2) \partial/\partial\varepsilon ((2t)^2 - \varepsilon^2)^{-1/2} + 7/360 (\pi/\beta)^4 \partial^3/\partial\varepsilon^3 ((2t)^2 - \varepsilon^2)^{-1/2} + 31/(21 \times 720) (\pi/\beta)^6 \partial^5/\partial\varepsilon^5 ((2t)^2 - \varepsilon^2)^{-1/2}] = n \quad (8)$$

where  $\varepsilon = \mu$  because of eq.(7).

Expression (8) can be developed up to the second order in  $\mu$  around the point  $\mu = 0$ , and then solved with the following result:

$$\mu = (\beta t)^6 (ns - 1) |t| / [1.1029 + .1694(\beta t)^2 + .0654(\beta t)^4] \quad (9)$$

In the case of weak hopping, this can be developed as

$$\mu = (\beta t)^6 (ns - 1) |t| [ .90671 - .13940(\beta t)^2 - .03284(\beta t)^4 ] \quad (9a)$$

It is obvious that  $\lim_{n,s \rightarrow 1} \mu = 0$ , which means that equations (9) and (9a) reduce to the solution discussed in [5].

A better approximation of the function  $\mu = \mu(\beta, t, n)$  can be obtained by pushing the development of eq. (8) around the point  $\mu = 0$  to higher orders. As a test, this development was performed to third order. However, solutions obtained in this way are too complicated for being applicable in calculations of the conductivity of Q1D organic conductors.

## Discussion

In this letter we have determined the dependence of the chemical potential of the electron gas on a one-dimensional lattice on the band-filling, hopping and temperature. The calculation was performed within the discrete Hubbard model. It was motivated by the need of introducing the doping in theoretical studies of the electrical conductivity of Q1D organic metals.

The development of eq. (8) in  $\mu$  around the point  $\mu = 0$  limits the applicability of eq. (9) to values of  $n$  not too different from 1. Work is already in progress aiming at an expression for the chemical potential which could be applied at arbitrary filling. Another line of progress concerns the application of the results of this letter to the analysis of the dependence of the conductivity of Q1D organic conductors on the doping.

## References

- [1] K.Bechgaard,C.S.Jacoben,K.Mortensen et al,Solid State Comm., **33**, 1119 (1980).
- [2] D.Jerome,A.Mazaud,M.Ribault and K.Bechgaard,J.Physique Lett., **41**, L95 (1980).
- [2a] J.P.Pouget,Mol.Cryst.Liq.Cryst., **230**, 101 (1993).
- [3] D.Jerome and F.Creuzet,in: Novel Superconductivity (ed.by S.A.Wolf and V.Z.Kresin), p.103 Plenum Press,New York (1987).
- [4] P.W.Anderson,Science, **235**, 1196 (1987).
- [5] E.H.Lieb and F.Y.Wu, Phys.Rev.Lett., **20**, 1445 (1968).
- [5a] V. Celebonović, poster AD2Po019 at the 14 EPS-CMD General Conference, Madrid ( Spain ) March 28.- 31.1994.
- [6] J. Voit,to appear in: Proc.of the NATO-ARW "The Physics and Mathematical Physics of the Hubbard Model", San Sebastian, ( Spain ), October 3 - 8, 1993, edited by D.K. Campbell, J.M.P. Carmelo and F. Guinea, Plenum Press, New York, (1994).
- [7] R. Kato, S. Aonuma, Y. Okano et al, Techn. Rep.of ISSP (Tokyo), **A2697**, 1 (1993). (to appear in Synth. Metals).
- [8] S. Fassari, J. Math. Phys., **30**, 1385 (1989).
- [9] R. P. Feynman, Statistical Mechanics: A Set of Lectures, W. A. Benjamin Inc., Reading, Mass.(1972).
- [10] E. T. Whittaker and G.N. Watson, A Course in Modern Analysis, Cambridge Univ. Press, Cambridge (1973).
- [11] V. K. Lukyanov, JINR Dubna preprint P4-93-300 (1993).
- [12] M. Abramowitz and I. A. Stegun, Handbook of Mathematical Functions, Dover Publ. Inc., New York (1972).
- [13] T. Dahm, L. Tewordt and S. Wermbter, Phys. Rev., **B49**, 748 (1994).
- [14] M. Imada, in: "Quantum Monte Carlo Methods in Condensed Matter Physics",pp.299-316, ed.by M.Suzuki,World Scientific,Singapore,(1993).
- [15] H.J.Rothe,Heidelberg preprint HD-THEP-93-12 and references given there (1993).